

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2004/030910

International filing date: 08 October 2004 (08.10.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/514,291
Filing date: 24 October 2003 (24.10.2003)

Date of receipt at the International Bureau: 21 May 2007 (21.05.2007)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

May 16, 2007

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/514,291

FILING DATE: October 24, 2003

RELATED PCT APPLICATION NUMBER: PCT/US04/30910

THE COUNTRY CODE AND NUMBER OF YOUR PRIORITY APPLICATION, TO BE USED FOR FILING ABROAD UNDER THE PARIS CONVENTION, IS US60/514,291




Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53 (c)


Docket Number		P-16348		Type a plus sign (+) inside this box -->	+
INVENTOR(s)/APPLICANT(s)					
LAST NAME	FIRST NAME	MIDDLE NAME	RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)		
GRAFF	JEREMY	RICHARD	INDIANAPOLIS, IN		
TITLE OF THE INVENTION (280 characters max)					
BISINDOLYL MALEIMIDES USEFUL FOR TREATING PROSTATE CANCER AND AKT-MEDIATED DISEASES					
CORRESPONDENCE ADDRESS					
Eli Lilly and Company Patent Division P.O. Box 6288 Indianapolis, Indiana 46206-6288			 25885 PATENT TRADEMARK OFFICE		
STATE	IN	ZIP CODE	46206-6288	COUNTRY	USA
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification	Number of pages	15	<input type="checkbox"/> Small Entity Statement		
<input type="checkbox"/> Drawing(s)	Number of Sheets		<input type="checkbox"/> Other (Specify)		
METHOD OF PAYMENT (check one)					
<input type="checkbox"/> A check or money order is enclosed to cover the Provisional filing fees				PROVISIONAL FILING FEE AMOUNT (\$)	\$160.00
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account Number:	05-0840				

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,
SIGNATURE



Date 10/23/03


TYPED or PRINTED NAME WILLIAM R. BOUDREAUX

REGISTRATION NO.
(if appropriate)

35,796

☐ Additional inventors are being named on separately numbered sheets attached hereto

PROVISIONAL APPLICATION FOR PATENT FILING ONLY

"Express Mail" mailing label number EL 832895863 US	Date of Deposit 10-24-03
I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to the Commissioner for Patents, P.O. Box 1450, Alexandria, VA, 22313-1450.	
Printed Name BUDEN THOMAS	Signature 

BISINDOLYL MALEIMIDES USEFUL FOR TREATING PROSTATE CANCER AND
AKT-MEDIATED DISEASES

BACKGROUND OF THE INVENTION

5

Protein kinases are involved in the signal transduction pathways linking growth factors, hormones and other cell regulation molecules to cell growth, survival and metabolism under both normal and pathological conditions. One such protein kinase, protein kinase B (also known as AKT), is a serine/threonine kinase that plays a central
10 role in promoting the proliferation and survival of a wide range of cell types, thereby protecting cells from apoptosis (programmed cell death) (Khwaja, *Nature* 33-34 (1990)). Three members of the AKT subfamily of second-messenger regulated serine/threonine protein kinases have been identified and are termed AKT-1, AKT-2, and AKT-3. A number of proteins involved in cell proliferation and survival have been described as
15 substrates of AKT in cells. Two examples of such substrates include glycogen synthase kinase-3 (GSK3) and Forkhead transcription factors (FKs). See Brazil and Hemmings, *Trends in Biochemical Sciences* 26, 675-664.

A number of protein kinases and phosphatases regulate the activity of AKT. For instance, activation of AKT is mediated by phosphatidylinositol 3-kinase (PI3-K), which
20 generates second messenger phospholipids that then bind to the pleckstrin homology (PH) binding domain of AKT. The binding attracts AKT to the plasma membrane where AKT is phosphorylated by phosphatidylinositol dependent kinase 1 (PDK1) at Thr308, which then triggers phosphorylation of AKT at Ser473 and activation of the enzyme. Amplifications of the catalytic subunit of PI3-K, p110 α , or mutations in the PI3-K
25 regulatory subunit, p85 α lead to activation of AKT in several types of human cancer. (Vivanco and Sawyers, *Nature Reviews in Cancer* (2002) 2: 489-501).

30

"Express Mail" mailing label number EL 832895863 US

Date of Deposit 10-24-03

35

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 C.F.R. 1.10 on the date indicated above and is addressed to the Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

QUEEN THOMAS
Printed Name

Queen Thomas
Signature

The tumor suppressor, PTEN, is a critical negative regulator of AKT activation by PI3-K (Myers *et al. Proc. Nat. Acad. Sci. 95, USA* (1998) 13513-13518). Inactivating mutations in the *Pten* gene have been found at high frequencies in a large number of human tumors and tumor cell lines, including prostate cancer, breast cancer, ovarian cancer, glioblastoma, melanoma and other cancer types. Inactivation of the PTEN protein results in elevated levels of phosphorylated AKT and increased AKT activity in tumor cells (Li, *et al., Science* (1997) 275: 1943-1947; Guldberg, *et al., Cancer Research* (1997) 57: 3660-3663; Risinger, *et al., Cancer Research* (1997) 57: 4736-4738; Vivanco and Sawyers, *Nature Reviews in Cancer* (2002) 2: 489-501). In addition to overactivation of AKT due to defects in PTEN, direct amplification and/or overexpression of AKT-2 and AKT-3 have been found in human neoplasia, for example ovarian, pancreatic, prostate and breast cancer cells (Cheung *et al., Proc. Nat. Acad. Sci. USA* (1992) 89:9267-9271; Cheung *et al., Proc. Nat. Acad. Sci. USA* (1996) 93:3636-3641; Nakatani *et al., J. Biol. Chem.* (1999) 274:21528-21532).


The critical role of AKT in cell proliferation and survival is further strengthened by studies showing that germline knockout of AKT-1 results in partial embryonic lethality. The surviving littermates display stunted growth, increased organismal apoptosis, and early deaths. (Cho *et al., J. Biol. Chem.* (2001) 276: 38349-38520; Chen *et al., Genes Dev.* (2001) 15: 2203-2208). It has also been demonstrated that pharmacological inactivation of AKT induces apoptosis in cultured human ovarian cancer cells (Yuan *et al., Oncogene* 19, 2324-2340, 2000) and decreases growth of a human ovarian carcinoma xenograft in mice (Hu *et al., Clin. Cancer Res.* 6, 880-886, 2000).

Recent studies have also demonstrated the role of the PI3-K/AKT pathway in the life cycle of numerous viruses. Some viral proteins have been shown to directly activate the PI3-K/AKT pathway, thus providing an environment favorable for viral replication. These include the Tat protein of human immunodeficiency virus (HIV), Protein X of hepatitis B virus, and NS5A of hepatitis C virus (Borgatti *et al., Eur. J. Immunol.* (1997) 27: 2805-2811; Lee *et al., J. Biol. Chem.* (2001) 276: 16969-16977; He *et al., J. Virol.* (2002) 76: 9207-9217). The PI3-K/AKT pathway is also required for initiation and completion of the replication cycle of human cytomegalovirus (HCMV). In fact, pharmacological inactivation of this pathway results in abortive production of HCMV and survival of the host cells (Johnson *et al., J. Virol.* (2001) 75: 6022-6032).

Because of its pivotal role in the regulation of cell survival, the PI3 kinase/ AKT pathway provides a novel therapeutic target for the effective treatment of various disorders, particularly cancer and viral infections. However, such treatment requires the development of potent, selective inhibitors of kinases within this pathway. Thus, the present invention provides a class of novel inhibitors of the PI3 kinase/AKT pathway, compositions comprising these compounds, and methods of using the compounds. Specifically, inhibition of PDK-1 by these compounds would be expected to suppress activation of the entire pathway as PDK-1 is the key kinase activating AKT. Inhibition of p70S6 kinase, a kinase effector downstream of AKT, would further suppress the enhanced ribosome biogenesis and protein translation triggered by AKT pathway activation. Prostatic adenocarcinoma (CaP) is the most common, non-cutaneous malignancy and the second-leading cause of cancer death in men. The disease has two distinct phases: the androgen-dependent phase, which can be treated effectively with androgen ablation therapies, and the androgen-independent phase. It is estimated that over thirty thousand men will die each year from androgen-independent metastatic CaP. Efforts to understand the metastatic progression of CaP progression to androgen-independent, metastatic disease involves a dampened apoptotic response, a release from the cell cycle block that initially follows androgen withdrawal and a shift from dependence on paracrine-derived growth and survival factors to autonomous production of these key proteins. Functional loss of the tumor suppressor phosphatase and tensin homologue deleted on the chromosome ten (PTEN) and subsequent activation of the AKT pathway, have been prominently implicated in the progression of CaP to androgen-independence. Activation of the AKT pathway can suppress the apoptotic response, undermine cell cycle control and selectively enhance the production of key growth and survival factors. Though many proteins and intracellular signaling pathways can influence these biological responses, activation of the AKT pathway is a particularly potent signal involved in CaP progression to androgen-independence and therefore provides a therapy of advanced androgen-independent CaP (Graff 2002). Treatment of CWR22Rv1, LNCaP and Du145 prostate cancer cells with the compound induces apoptosis.

SUMMARY OF THE INVENTION

The present invention provides a method of treating prostate cancer comprising administering to a patient in need thereof a therapeutically effective amount of a
5 compound of the formula (I)



(I)

wherein R¹ and R² are each independently hydrogen or C₁-C₄ alkyl; or a pharmaceutically acceptable salt thereof.

10 In a second embodiment, the invention provides a method of treating androgen-independent prostatic adenocarcinoma comprising administering to a patient in need thereof a therapeutically effective amount of compound of formula (I) or a pharmaceutically acceptable salt thereof.

15 In a third embodiment, the invention provides a method of treating an AKT-mediated disease selected from the group consisting of glioblastoma, colon cancer, pancreatic cancer, ovarian cancer, endometrial cancer, and renal cell cancer, comprising administering to a patient in need thereof a therapeutically effective amount of compound of formula (I) or a pharmaceutically acceptable salt thereof.

DETAILED DESCRIPTION OF THE INVENTION

General terms used in the description of compounds herein described bear their usual meanings. For example, the term "C₁-C₄ alkyl" refers to straight or branched, monovalent, saturated aliphatic chains of 1 to 4 carbon atoms and includes, but is not
5 limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, *sec*-butyl, and *tert*-butyl.

Preferred compounds of this invention include compounds of formula I wherein R¹ is hydrogen, methyl, ethyl, n-propyl, or isopropyl. Further preferred compounds include those wherein R² is hydrogen or methyl. More preferred compounds are those where R¹ is hydrogen. The skilled artisan will appreciate that additional preferred
10 embodiments may be selected by combining the preferred embodiments above, or by reference to the examples given herein.

The term "pharmaceutically-acceptable salt" as used herein, refers to a salt of a compound of the above Formula (I). It should be recognized that the particular counterion forming a part of any salt of this invention is usually not of a critical nature, so
15 long as the salt as a whole is pharmacologically acceptable and as long as the counterion does not contribute undesired qualities to the salt as a whole.

The compounds of Formula (I) described herein form pharmaceutically-acceptable acid addition salts with a wide variety of organic and inorganic acids and include the physiologically-acceptable salts which are often used in pharmaceutical chemistry. Such
20 salts are also part of this invention. A pharmaceutically-acceptable acid addition salt is formed from a pharmaceutically-acceptable acid, as is well known in the art. Such salts include the pharmaceutically acceptable salts listed in Journal of Pharmaceutical Science, 66, 2-19 (1977), which are known to the skilled artisan. *See also*, The Handbook of Pharmaceutical Salts; Properties, Selection, and Use. P. H. Stahl and C. G. Wermuth
25 (ED.s), Verlag, Zurich (Switzerland) 2002.

Typical inorganic acids used to form such salts include hydrochloric, hydrobromic, hydriodic, nitric, sulfuric, phosphoric, hypophosphoric, metaphosphoric, pyrophosphoric, and the like. Salts derived from organic acids, such as aliphatic mono and dicarboxylic acids, phenyl substituted alkanoic acids, hydroxyalkanoic and
30 hydroxyalkandioic acids, aromatic acids, aliphatic and aromatic sulfonic acids, may also be used. Such pharmaceutically acceptable salts thus include acetate, phenylacetate, trifluoroacetate, acrylate, ascorbate, benzoate, chlorobenzoate, dinitrobenzoate,

hydroxybenzoate, methoxybenzoate, methylbenzoate, o-acetoxybenzoate, naphthalene-2-benzoate, bromide, isobutyrate, phenylbutyrate, α -hydroxybutyrate, butyne-1,4-dicarboxylate, hexyne-1,4-dicarboxylate, caprate, caprylate, cinnamate, citrate, formate, fumarate, glycollate, heptanoate, hippurate, lactate, malate, maleate, hydroxymaleate, 5 malonate, mandelate, mesylate, nicotinate, isonicotinate, nitrate, oxalate, phthalate, teraphthalate, propiolate, propionate, phenylpropionate, salicylate, sebacate, succinate, suberate, benzenesulfonate, p-bromobenzenesulfonate, chlorobenzenesulfonate, ethylsulfonate, 2-hydroxyethylsulfonate, methylsulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, naphthalene-1,5-sulfonate, p-toluenesulfonate, xylenesulfonate, 10 tartarate, and the like.

The compounds of formula (I) are described in Heath, Jr. et al., U.S. Pat. No. 5,668,152. The synthesis of the compounds of formula (I) are fully set forth as well as a disclosure that said compounds are useful as beta-1 and beta-2 isozyme selective protein kinase C (PKC) inhibitors. As isozyme selective PKC inhibitors, the compounds have 15 previously been disclosed as useful in the treatment of conditions associated with diabetes mellitus and its complications as well ischemia, inflammation, central nervous system disorders, cardiovascular disease, dermatological disease, Alzheimer's disease and cancer. U.S. Pat. No. 5,668,152 is hereby incorporated by reference in its entirety as if fully set forth.

20 While U.S. Pat. No. 5,668,152 describes the treatment of cancer using PKC beta-1 and beta-2 selective inhibitors, of which the present compounds of formula (I) are included generically, there is no teaching or suggestion that the compounds of formula (I) are inhibitors of the PI3K/ AKT pathway. Because the AKT pathway acts as a central regulator of the apoptotic response, inhibitors of this pathway would be expected to 25 induce apoptosis and/ or block cell cycle progression whereas inhibition of PKC, which has many disparate roles in the cell, would not necessarily be expected to do so.

As used herein, the term "patient" refers to a warm-blooded animal or mammal which is in need of treating, or at risk of developing, one or more diseases or disorders associated with AKT pathway activity (e.g. PDK-1/ p70S6 kinase activity). It is 30 understood that guinea pigs, dogs, cats, rats, mice, hamsters, and primates, including humans, are examples of patients within the scope of the meaning of the term. Preferred patients include humans.

The compounds of the present invention can be administered alone or in the form of a pharmaceutical composition, that is, combined with pharmaceutically acceptable carriers, or excipients, the proportion and nature of which are determined by the solubility and chemical properties of the compound selected, the chosen route of administration, and standard pharmaceutical practice. The compounds of the present invention, while effective themselves, may be formulated and administered in the form of their pharmaceutically acceptable salts, for purposes of stability, convenience of crystallization, increased solubility, and the like.

Thus, the present invention provides pharmaceutical compositions comprising a compound of the Formula (I) and a pharmaceutically acceptable diluent.

The compounds of Formula (I) can be administered by a variety of routes. In effecting treatment of a patient afflicted with or at risk of developing the disorders described herein, a compound of Formula (I) can be administered in any form or mode that makes the compound bioavailable in an effective amount, including oral and parenteral routes. For example, compounds of Formula (I) can be administered orally, by inhalation, or by the subcutaneous, intramuscular, intravenous, transdermal, intranasal, rectal, ocular, topical, sublingual, buccal, or other routes. Oral administration is generally preferred for treatment of the disorders described herein. However, oral administration is not the only preferred route. For example, the intravenous route may be preferred as a matter of convenience or to avoid potential complications related to oral administration. When the compound of Formula (I) is administered through the intravenous route, an intravenous bolus or slow infusion is preferred.

One skilled in the art of preparing formulations can readily select the proper form and mode of administration depending upon the particular characteristics of the compound selected, the disorder or condition to be treated, the stage of the disorder or condition, and other relevant circumstances. (*Remington's Pharmaceutical Sciences*, 18th Edition, Mack Publishing Co. (1990)).

The pharmaceutical compositions are prepared in a manner well known in the pharmaceutical art. The carrier or excipient may be a solid, semi-solid, or liquid material that can serve as a vehicle or medium for the active ingredient. Suitable carriers or excipients are well known in the art. The pharmaceutical composition may be adapted for oral, inhalation, parenteral, or topical use and may be administered to the patient in the

form of tablets, capsules, aerosols, inhalants, suppositories, solutions, suspensions, or the like.

For the purpose of oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of tablets, troches, capsules, elixirs, suspensions, syrups, wafers, chewing gums and the like. These preparations should contain at least 4% of the compound of the present invention, the active ingredient, but may be varied depending upon the particular form and may conveniently be between 4% to about 70% of the weight of the unit. The amount of the compound present in compositions is such that a suitable dosage will be obtained. Preferred compositions and preparations according to the present invention may be determined by a person skilled in the art.

The tablets, pills, capsules, troches, and the like may also contain one or more of the following adjuvants: binders such as povidone, hydroxypropyl cellulose, microcrystalline cellulose, gum tragacanth or gelatin; excipients such as dicalcium phosphate, starch, or lactose; disintegrating agents such as alginic acid, Primogel, corn starch and the like; lubricants such as talc, hydrogenated vegetable oil, magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; and sweetening agents, such as sucrose, aspartame, or saccharin, or a flavoring agent, such as peppermint, methyl salicylate or orange flavoring, may be added. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or a fatty oil. Other dosage unit forms may contain other various materials that modify the physical form of the dosage unit, for example, coatings. Thus, tablets or pills may be coated with sugar, shellac, or other coating agents. Syrups may contain, in addition to the present compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors. Materials used in preparing these various compositions should be pharmaceutically pure and non-toxic in the amounts used.

The compounds of Formula (I) are inhibitors of PDK1 and p70S6 kinase, two members of the PI3kinase/ AKT pathway. The inhibitory activity of the compounds of Formula (I) may be demonstrated by the methods below.

Kinase Activity Assays

The assay described measures the phosphorylation of the PDK1 consensus phosphorylation site PDK-tide peptide


(KTFCGTPEYLAPEVRREPRILSEEEQEMFRDFDYIADWC; cat # 14-452, lot

- 5 23876U) by recombinant PDK-1 (UBI) at Km for ATP and PDKtide saturation using phosphocellulose membrane filter plates. Phosphorylation of the p70S6 kinase substrate by recombinant p70S6 kinase is also measured similarly.

Cell culture, Drug Treatment, Apoptosis and Proliferation Assays

- 10 Both HCT116 colon carcinoma (cat#CCL-247) and U87MG glioblastoma (cat#HTB-14) cell lines were obtained from the American Type Culture Collection (ATCC). The standard growth media differed for each cell line but all were grown in 10% heat-inactivated FBS (Invitrogen cat# 10082-147), 37°C, 5% CO₂ atmosphere and in a humidified chamber. Cell passage was completed one to two times per week using
- 15 0.25% trypsin / 1mM EDTA (Invitrogen,cat# 25200-056) solution maintaining cells in log phase growth. U87MG cells were cultured in DMEM media (Invitrogen cat# 11965-092), 1mM non-essential amino acids (NEAA), and 0.1mM Sodium Pyruvate. HCT116 cells were grown in McCoy's 5A Modified media (Invitrogen, cat# 16600-082), 0.15% sodium bicarbonate, 0.1mM HEPES, 25mM D-glucose and 0.1mM sodium Pyruvate.
- 20 Apoptosis assays were executed using the Cell Death Detection ELISA^{plus} (Roche, 1774425) assay kit strictly following the enclosed protocol.

Changes in cellular proliferation resulting from treatment with LY317615, which is a compound of the formula



LY317615

5 or a compound of formula (I) where R^1 is hydrogen and R^2 is methyl (Compound 1) were assessed by incorporation of propidium iodide (PI) (Sigma, cat# p-4864). Briefly, each cell culture plate was centrifuged 10 minutes (200rpm), the supernatant was gently aspirated and 100 μ l 0.125 mM PI in PBS was added to each well of a 96-well plate. The fluorescence intensity of each well in the culture was measured (non-viable cells) using the Vector² multi-channel plate reader (Wallac, model#1420) and frozen to -80° C. The plate was allowed to thaw, come to room temperature and re-analyzed for changes in fluorescence intensity (total cells) again using the Vector². The proliferating cells in the culture were determined by subtracting the non-viable fraction from the total cells. The results were then reported as a percent of the un-treated control.

15 Protein lysates were prepared by incubation in RIPA Buffer (50 mM Tris -HCl, 150 mM NaCl, 1 mM EDTA, 1% NP-40, 0.25% Sodium deoxycholate, 1 mM sodium fluoride, 1mM sodium orthovanadate and CompleteTM protease inhibitors (Roche Cat# 10019600) for one hour with rotation. The lysate was then centrifuged (10 minutes@ 10,000 rpm), supernatant harvested and protein concentrations were determined using the Bio-Rad DC Protein Assay (cat# 500-0122). Proteins were separated by SDS-PAGE using 4-20% tris-glycine gels (Invitrogen, cat# EC6028) and transferred to HyperTM-bond PVDF membrane (Amersham, cat# rpn303F). All primary antibodies were incubated overnight at 4° C in 5% milk / 1X PBS (Gibco, cat# 70011-044) solution. Horseradish peroxidase (HRP) linked secondary antibodies (Santa Cruz, cat# sc-2055, sc-2054) were

incubated for a minimum of two hours prior to detection. Specific signal was determined by the Lumi-Imager™ and Lumi-Analyst software to define changes in protein expression and phosphorylation. The primary antibodies used are as follows: GSK3b, pGSK3b^{ser9} (Cell Signaling, cat#9332, 9336), S6 ribosomal protein, pS6 ribosomal protein^{ser240/244} (Cell Signaling, cat#2212, 2215), AKT (Transduction Labs cat#610861), pAKT^{ser308}, pAKT^{ser473} (Cell Signaling, cat#9275, 9271), PHASi (Zymed, cat#51-2900), p4EBP1^{ser65} (Cell Signaling cat#9451), p70S6 kinase, phos-p70S6 kinase^{thr421/ser424} (Cell Signaling cat#9202, 9204), p90RSK, phos-p90RSK^{thr359/ser363} (Cell Signaling cat#9347, 9344), FKHL1, pFKHL1^{thr32} (Upstate Biochemicals, Inc. cat#06-951, 06952).

Experimental Protocol for *In Vivo* Tumor Inhibition Studies


Approximately 5 x10⁶ tumor cells are implanted subcutaneously into the flank of athymic nude mice (Harlan, Indianapolis, IN). Treatment of tumors begins when the tumors reach 100mm³ and continues for 21 consecutive days twice per day PO. Body weight and tumor size are monitored weekly or twice weekly.

Results

Compound 1 inhibits PDK-1 with an EC50 of 370 nM and inhibits p70S6kinase with an EC50 <500nM. Treatment of human cancer cell lines with Compound 1 induces apoptosis and suppresses phosphorylation of GSK3β, the forkhead transcription factor AFX, 4EBP1, and ribosomal protein- all readouts of AKT pathway activity. Furthermore, treatment of HCT116 human tumor xenograft-bearing mice with Compound 1 suppresses GSK3β ser9 phosphorylation for up to 8 hours post dosing. Anti-tumor efficacy of the compound has been demonstrated in both HCT116 colon cancer xenografts and in U87MG glioblastoma xenografts (p < 0.01 for each study).

WHAT IS CLAIMED IS:

1. A method of treating prostate cancer comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the formula (I)




(I)

wherein R¹ and R² are each independently hydrogen or C₁-C₄ alkyl; or a pharmaceutically acceptable salt thereof.

2. A method according to claim 1 wherein R² is hydrogen or methyl, or a pharmaceutically acceptable salt thereof.
3. A method according to claim 2 wherein R¹ is hydrogen, methyl, ethyl, n-propyl, or isopropyl, or a pharmaceutically acceptable salt thereof.
4. A method according to claim 1 wherein R¹ is hydrogen and R² is methyl, or a pharmaceutically acceptable salt thereof.
5. A method according to claim 1 wherein said patient is a human diagnosed with prostate cancer.
6. A method according to claim 1 wherein said patient is a human at risk of developing prostate cancer.

7. A method of treating androgen-independent prostatic adenocarcinoma comprising administering to a patient in need thereof a therapeutically effective amount of a compound of the formula (I)



(I)

wherein R¹ and R² are each independently hydrogen or C₁-C₄ alkyl; or a pharmaceutically acceptable salt thereof.

8. A method according to claim 7 wherein R² is hydrogen or methyl, or a pharmaceutically acceptable salt thereof.


9. A method according to claim 8 wherein R¹ is hydrogen, methyl, ethyl, n-propyl, or isopropyl, or a pharmaceutically acceptable salt thereof.

10. A method according to claim 7 wherein R¹ is hydrogen and R² is methyl, or a pharmaceutically acceptable salt thereof.

11. A method according to claim 7 wherein said patient is a human diagnosed with androgen-independent prostatic adenocarcinoma.

12. A method according to claim 7 wherein said patient is a human at risk of developing androgen-independent prostatic adenocarcinoma.

13. A method according to treating an AKT-mediated disease selected from the group consisting of glioblastoma, colon cancer, pancreatic cancer, ovarian cancer, endometrial cancer, and renal cell cancer, comprising administering to a patient in need thereof a therapeutically effective amount of compound of formula (I)



(I)

wherein R^1 and R^2 are each independently hydrogen or C_1 - C_4 alkyl; or a pharmaceutically acceptable salt thereof.

14. A method according to claim 13 wherein said AKT-mediated disease is glioblastoma.

15. A method according to claim 13 wherein said AKT-mediated disease is colon cancer.

16. A method according to claim 13 wherein said AKT-mediated disease is pancreatic cancer.

17. A method according to claim 13 wherein said AKT-mediated disease is ovarian cancer.


18. A method according to claim 13 wherein said AKT-mediated disease is endometrial cancer.

19. A method according to claim 13 wherein said AKT-mediated disease is renal cell cancer.

20. A method according to claim 13 wherein R^1 is hydrogen and R^2 is methyl, or a pharmaceutically acceptable salt thereof.

ABSTRACT

The present invention provides a method of treating prostate cancer comprising administering to a patient in need thereof a therapeutically effective amount of a
5 compound of the formula (I)



(I)

wherein R¹ and R² are each independently hydrogen or C₁-C₄ alkyl; or a pharmaceutically acceptable salt thereof.

10 In a second embodiment, the invention provides a method of treating androgen-independent prostatic adenocarcinoma comprising administering to a patient in need thereof a therapeutically effective amount of compound of formula (I) or a pharmaceutically acceptable salt thereof.

15 In a third embodiment, the invention provides a method of treating an AKT-mediated disease selected from the group consisting of glioblastoma, colon cancer, pancreatic cancer, ovarian cancer, endometrial cancer, and renal cell cancer, comprising administering to a patient in need thereof a therapeutically effective amount of compound of formula (I) or a pharmaceutically acceptable salt thereof.